


What is claimed is:

[illegible]

1. A compound of formula



wherein W^3 and X^3 are independently selected from the group consisting of $-CR^1R^2$, $-O-$, $-NR^3$, $-S-$, and $-Se$; Y^3 is selected from the group consisting of $-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$,

5 $-(CH_2)_a-NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$,

$-(CH_2)_a-N(R^3)-(CH_2)_b-CONH-Bm$, $(CH_2)_a-N(R^3)-(CH_2)_c-NHCO-Bm$,

$-(CH_2)_a-N(R^3)-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$,

$-(CH_2)_a-N(R^3)-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$,

$-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-(CH_2)_a-CONH-Bm$,

10 $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-(CH_2)_a-NHCO-Bm$,

$-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-CH_2-(CH_2OCH_2)_d-CONH-Bm$,

$-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-CH_2-(CH_2OCH_2)_d-NHCO-Bm$, $-(CH_2)_a-NR^3R^4$,

and $-CH_2(CH_2OCH_2)_b-CH_2NR^3R^4$; Z^3 is selected from the group consisting of $-(CH_2)_a-CONH-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Dm$, $-(CH_2)_a-NHCO-Dm$,

15 $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Dm$, $-(CH_2)_a-N(R^3)-(CH_2)_b-CONH-Dm$,

$(CH_2)_a-N(R^3)-(CH_2)_c-NHCO-Dm$, $-(CH_2)_a-N(R^3)-CH_2-(CH_2OCH_2)_b-CH_2-CONH-$
 Dm , $-(CH_2)_a-N(R^3)-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-$
 $N(R^3)-(CH_2)_a-CONH-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-(CH_2)_a-NHCO-Dm$,
5 $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-CH_2-(CH_2OCH_2)_d-CONH-Dm$, $-CH_2-(CH_2OCH_2)_b-$
 $CH_2-N(R^3)-CH_2-(CH_2OCH_2)_d-NHCO-Dm$, $-(CH_2)_a-NR^3R^4$, and
 $-CH_2(CH_2OCH_2)_b-CH_2NR^3R^4$; A_1 is a single or a double bond; B_1 , C_1 , and D_1
20 are independently selected from the group consisting of $-O-$, $-S-$, $-Se-$, $-P-$,
 $-CR^1R^2$, $-CR^1$, alkyl, NR^3 , and $-C=O$; A_1 , B_1 , C_1 , and D_1 may together form
a 6- to 12-membered carbocyclic ring or a 6- to 12-membered
heterocyclic ring optionally containing one or more oxygen, nitrogen, or
sulfur atom; a_3 and b_3 are independently from 0 to 5; R^1 to R^4 , and R^{29} to
25 R^{37} are independently selected from the group consisting of hydrogen,
 C_1-C_{10} alkyl, C_5-C_{20} aryl, C_1-C_{10} alkoxy, C_1-C_{10} polyalkoxyalkyl, C_1-C_{20}
polyhydroxyalkyl, C_5-C_{20} polyhydroxyaryl, C_1-C_{10} aminoalkyl, cyano, nitro,
halogen, saccharide, peptide, $-CH_2(CH_2OCH_2)_b-CH_2-OH$, $-(CH_2)_a-CO_2H$,
 $-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$, $-(CH_2)_a-NHCO-Bm$,
30 $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$, $-(CH_2)_a-OH$ and $-CH_2-(CH_2OCH_2)_b-CO_2H$;
 Bm and Dm are independently selected from the group consisting of a
bioactive peptide, a protein, a cell, an antibody, an antibody fragment, a
saccharide, a glycopeptide, a peptidomimetic, a drug, a drug mimic, a
hormone, a metal chelating agent, a radioactive or nonradioactive metal
35 complex, and an echogenic agent; a and c are independently from 1 to
20; and b and d are independently vary from 1 to 100.

2. The compound of claim 1 wherein W^3 and X^3 are independently selected from the group consisting of $-C(CH_3)_2$, $-C((CH_2)_aOH)CH_3$, $-C((CH_2)_aOH)_2$, $-C((CH_2)_aCO_2H)CH_3$, $-C((CH_2)_aCO_2H)_2$, $-C((CH_2)_aNH_2)CH_3$, $C((CH_2)_aNH_2)_2$, $C((CH_2)_aNR^3R^4)_2$, $-NR^3$, and $-S-$; Y^3 is selected from the group consisting of $-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$, $-(CH_2)_a-NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$, $-(CH_2)_a-NR^3R^4$, and $-CH_2(CH_2OCH_2)_b-CH_2NR^3R^4$; Z^3 is selected from the group consisting of $-(CH_2)_a-CONH-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Dm$, $-(CH_2)_a-NHCO-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Dm$, $-(CH_2)_a-NR^3R^4$, and $-CH_2(CH_2OCH_2)_b-CH_2NR^3R^4$; A_1 is a single or a double bond; B_1 , C_1 , and D_1 are independently selected from the group consisting of $-O-$, $-S-$, NR^3 , $(CH_2)_a-CR^1R^2$, and $-CR^1$; A_1 , B_1 , C_1 , and D_1 may together form a 6- to 10-membered carbocyclic ring or a 6- to 10-membered heterocyclic ring optionally containing one or more oxygen, nitrogen, or sulfur atom; a_3 and b_3 are independently from 0 to 3; R^1 to R^4 , and R^{29} to R^{37} are independently selected from the group consisting of hydrogen, C_1 - C_{10} alkyl, C_5 - C_{12} aryl, C_1 - C_{10} alkoxy, C_1 - C_{10} polyhydroxyalkyl, C_5 - C_{12} polyhydroxyaryl, C_1 - C_{10} aminoalkyl, mono- or oligosaccharide, peptide with 2 to 30 amino acid units, $-CH_2(CH_2OCH_2)_b-CH_2-OH$, $-(CH_2)_a-CO_2H$, $-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$, $-(CH_2)_a-NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$, $-(CH_2)_a-OH$ and $-CH_2-(CH_2OCH_2)_b-CO_2H$; Bm and Dm are independently selected from the group consisting of a bioactive peptide containing 2 to 30 amino acid


25

5

4. A method for performing a diagnostic or therapeutic procedure comprising

administering to an individual an effective amount of the

compound of formula



- 5 wherein W^3 and X^3 are independently selected from the group consisting of $-CR^1R^2$, $-O-$, $-NR^3$, $-S-$, and $-Se$; Y^3 is selected from the group consisting of $-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$, $-(CH_2)_a-NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$, $-(CH_2)_a-N(R^3)-(CH_2)_b-CONH-Bm$, $(CH_2)_a-N(R^3)-(CH_2)_c-NHCO-Bm$, $-(CH_2)_a-N(R^3)-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$, $-(CH_2)_a-N(R^3)-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-(CH_2)_a-NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-CH_2-(CH_2OCH_2)_d-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-CH_2-(CH_2OCH_2)_d-NHCO-Bm$, $-(CH_2)_a-NR^3R^4$, and $-CH_2(CH_2OCH_2)_b-CH_2NR^3R^4$; Z^3 is selected from the group consisting of $-(CH_2)_a-CONH-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Dm$, $-(CH_2)_a-NHCO-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Dm$, $-(CH_2)_a-N(R^3)-(CH_2)_b-CONH-Dm$, $(CH_2)_a-N(R^3)-(CH_2)_c-NHCO-Dm$,
- 10
- 15

- $-(CH_2)_a-N(R^3)-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Dm$, $-(CH_2)_a-N(R^3)-CH_2-$
 $(CH_2OCH_2)_b-CH_2-NHCO-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-(CH_2)_a-CONH-$
 Dm , $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-(CH_2)_a-NHCO-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-$
20 $N(R^3)-CH_2-(CH_2OCH_2)_d-CONH-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-CH_2-$
 $(CH_2OCH_2)_d-NHCO-Dm$, $-(CH_2)_a-NR^3R^4$, and $-CH_2(CH_2OCH_2)_b-CH_2NR^3R^4$; A_1
is a single or a double bond; B_1 , C_1 , and D_1 are independently selected
from the group consisting of $-O-$, $-S-$, $-Se-$, $-P-$, $-CR^1R^2$, $-CR^1$, alkyl, NR^3 ,
and $-C=O$; A_1 , B_1 , C_1 , and D_1 may together form a 6- to 12-membered
25 carbocyclic ring or a 6- to 12-membered heterocyclic ring optionally
containing one or more oxygen, nitrogen, or sulfur atom; a_3 and b_3 are
independently from 0 to 5; R^1 to R^4 , and R^{29} to R^{37} are independently
selected from the group consisting of hydrogen, C_1 - C_{10} alkyl, C_5 - C_{20} aryl,
 C_1 - C_{10} alkoxy, C_1 - C_{10} polyalkoxyalkyl, C_1 - C_{20} polyhydroxyalkyl, C_5 - C_{20}
30 polyhydroxyaryl, C_1 - C_{10} aminoalkyl, cyano, nitro, halogen, saccharide,
peptide, $-CH_2(CH_2OCH_2)_b-CH_2-OH$, $-(CH_2)_a-CO_2H$,
 $-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$, $-(CH_2)_a-NHCO-Bm$,
 $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$, $-(CH_2)_a-OH$ and $-CH_2-(CH_2OCH_2)_b-CO_2H$;
 Bm and Dm are independently selected from the group consisting of a
35 bioactive peptide, a protein, a cell, an antibody, an antibody fragment, a
saccharide, a glycopeptide, a peptidomimetic, a drug, a drug mimic, a
hormone, a metal chelating agent, a radioactive or nonradioactive metal
complex, and an echogenic agent; a and c are independently from 1 to

20; and b and d are independently from 1 to 100, and a pharmaceutically
40 acceptable carrier or excipient to form a composition,
activating the compound using light, and
performing the diagnostic or therapeutic procedure.

5. The method of claim 4 comprising administering to an individual an effective amount of the compound wherein W^3 and X^3 are independently selected from the group consisting of $-C(CH_3)_2$, $-C((CH_2)_aOH)CH_3$, $-C((CH_2)_aOH)_2$, $-C((CH_2)_aCO_2H)CH_3$, $-C((CH_2)_aCO_2H)_2$,
5 $-C((CH_2)_aNH_2)CH_3$, $C((CH_2)_aNH_2)_2$, $C((CH_2)_aNR^3R^4)_2$, $-NR^3$, and $-S-$; Y^3 is selected from the group consisting of $-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$, $-(CH_2)_a-NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$, $-(CH_2)_a-NR^3R^4$, and $-CH_2(CH_2OCH_2)_b-CH_2NR^3R^4$; Z^3 is selected from the group consisting of $-(CH_2)_a-CONH-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Dm$, $-(CH_2)_a-NHCO-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Dm$, $-(CH_2)_a-NR^3R^4$, and $-CH_2(CH_2OCH_2)_b-CH_2NR^3R^4$; A_1 is a single or a double bond;
10 B_1 , C_1 , and D_1 are independently selected from the group consisting of $-O-$, $-S-$, NR^3 , $(CH_2)_a-CR^1R^2$, and $-CR^1$; A_1 , B_1 , C_1 , and D_1 may together form a 6- to 10-membered carbocyclic ring or a 6- to 10-membered
15 heterocyclic ring optionally containing one or more oxygen, nitrogen, or sulfur atom; a_3 and b_3 independently vary from 0 to 3; R^1 to R^4 , and R^{29} to R^{37} are independently selected from the group consisting of hydrogen, C_1 - C_{10} alkyl, C_5 - C_{12} aryl, C_1 - C_{10} alkoxy, C_1 - C_{10} polyhydroxyalkyl, C_5 - C_{12}

polyhydroxyaryl, C₁-C₁₀ aminoalkyl, mono- or oligosaccharide, peptide with 2 to 30 amino acid units, -CH₂(CH₂OCH₂)_b-CH₂-OH, -(CH₂)_a-CO₂H, -(CH₂)_a-CONH-Bm, -CH₂-(CH₂OCH₂)_b-CH₂-CONH-Bm, -(CH₂)_a-NHCO-Bm, -CH₂-(CH₂OCH₂)_b-CH₂-NHCO-Bm, -(CH₂)_a-OH and -CH₂-(CH₂OCH₂)_b-CO₂H; Bm and Dm are independently selected from the group consisting of a bioactive peptide containing 2 to 30 amino acid units, an antibody, a
25 mono- or oligosaccharide, a glycopeptide, a metal chelating agent, a radioactive or nonradioactive metal complex, and an echogenic agent; a and c are independently from 1 to 10; and b and d are independently from 1 to 30.

6. The method of claim 5 comprising administering to an individual an effective amount of the compound wherein each of W³ and X³ is C((CH₂)OH)₂; Y³ is -(CH₂)₂-CONH-Bm; Z³ is -(CH₂)₂-CONH-Dm; A₁ is a single bond; A₁, B₁, C₁, and D₁ together form a 6-membered carbocyclic
5 ring; each a₃ and b₃ is 1; R²⁹ is galactose; each R³⁰ to R³⁷ is hydrogen; Bm is Octreotate; and Dm is bombesin (7-14).

7. The method of claim 4 wherein said procedure uses light of wavelength in the region of 350-1300 nm.

8. The method of claim 4 wherein the diagnostic procedure is optical tomography.

9. The method of claim 4 wherein the diagnostic procedure is fluorescence endoscopy.

10. The method of claim 4 further comprising monitoring a blood clearance profile of said compound by a method selected from the group consisting of fluorescence, absorbance, and light scattering, wherein light of wavelength in the region of 350-1300 nm is used.


11. The method of claim 4 wherein said procedure further comprises imaging and therapy, wherein said imaging and therapy is selected from the group consisting of absorption, light scattering, photoacoustic and sonofluoresence technique.

12. The method of claim 4 wherein said procedure is capable of diagnosing atherosclerotic plaques and blood clots.

13. The method of claim 4 wherein said procedure comprises administering localized therapy.

14. The method of claim 4 wherein said therapeutic procedure comprises photodynamic therapy.

18. A composition comprising a cyanine dye bioconjugate of formula



- wherein W^3 and X^3 are independently selected from the group consisting of $-CR^1R^2$, $-O-$, $-NR^3$, $-S-$, and $-Se$; Y^3 is selected from the group consisting of $-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$, $-(CH_2)_a-$
- 5 $NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$, $-(CH_2)_a-N(R^3)-(CH_2)_b-CONH-Bm$, $(CH_2)_a-N(R^3)-(CH_2)_c-NHCO-Bm$, $-(CH_2)_a-N(R^3)-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$, $-(CH_2)_a-N(R^3)-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-(CH_2)_a-NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-CH_2-(CH_2OCH_2)_d-CONH-Bm$,
- 10 $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-CH_2-(CH_2OCH_2)_d-NHCO-Bm$, $-(CH_2)_a-NR^3R^4$, and $-CH_2(CH_2OCH_2)_b-CH_2NR^3R^4$; Z^3 is selected from the group consisting of $-(CH_2)_a-CONH-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Dm$, $-(CH_2)_a-NHCO-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Dm$, $-(CH_2)_a-N(R^3)-(CH_2)_b-CONH-Dm$, $(CH_2)_a-N(R^3)-(CH_2)_c-NHCO-Dm$, $-(CH_2)_a-N(R^3)-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Dm$,
- 15 $-(CH_2)_a-N(R^3)-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-(CH_2)_a-CONH-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-N(R^3)-(CH_2)_a-NHCO-Dm$,

$-\text{CH}_2-(\text{CH}_2\text{OCH}_2)_b-\text{CH}_2-\text{N}(\text{R}^3)-\text{CH}_2-(\text{CH}_2\text{OCH}_2)_d-\text{CONH-Dm}$, $-\text{CH}_2-(\text{CH}_2\text{OCH}_2)_b-$
 $\text{CH}_2-\text{N}(\text{R}^3)-\text{CH}_2-(\text{CH}_2\text{OCH}_2)_d-\text{NHCO-Dm}$, $-(\text{CH}_2)_a-\text{NR}^3\text{R}^4$, and
 $-\text{CH}_2(\text{CH}_2\text{OCH}_2)_b-\text{CH}_2\text{NR}^3\text{R}^4$; A_1 is a single or a double bond; B_1 , C_1 , and D_1
20 are independently selected from the group consisting of $-\text{O}-$, $-\text{S}-$, $-\text{Se}-$, $-\text{P}-$,
 $-\text{CR}^1\text{R}^2$, $-\text{CR}^1$, alkyl, NR^3 , and $-\text{C}=\text{O}$; A_1 , B_1 , C_1 , and D_1 may together form
a 6- to 12-membered carbocyclic ring or a 6- to 12-membered
heterocyclic ring optionally containing one or more oxygen, nitrogen, or
sulfur atom; a_3 and b_3 are independently from 0 to 5; R^1 to R^4 , and R^{29} to
25 R^{37} are independently selected from the group consisting of hydrogen, C_1 -
 C_{10} alkyl, C_5 - C_{20} aryl, C_1 - C_{10} alkoxy, C_1 - C_{10} polyalkoxyalkyl, C_1 - C_{20}
polyhydroxyalkyl, C_5 - C_{20} polyhydroxyaryl, C_1 - C_{10} aminoalkyl, cyano, nitro,
halogen, saccharide, peptide, $-\text{CH}_2(\text{CH}_2\text{OCH}_2)_b-\text{CH}_2-\text{OH}$, $-(\text{CH}_2)_a-\text{CO}_2\text{H}$,
 $-(\text{CH}_2)_a-\text{CONH-Bm}$, $-\text{CH}_2-(\text{CH}_2\text{OCH}_2)_b-\text{CH}_2-\text{CONH-Bm}$, $-(\text{CH}_2)_a-\text{NHCO-Bm}$,
30 $-\text{CH}_2-(\text{CH}_2\text{OCH}_2)_b-\text{CH}_2-\text{NHCO-Bm}$, $-(\text{CH}_2)_a-\text{OH}$ and $-\text{CH}_2-(\text{CH}_2\text{OCH}_2)_b-\text{CO}_2\text{H}$;
 Bm and Dm are independently selected from the group consisting of a
bioactive peptide, a protein, a cell, an antibody, an antibody fragment, a
saccharide, a glycopeptide, a peptidomimetic, a drug, a drug mimic, a
hormone, a metal chelating agent, a radioactive or nonradioactive metal
35 complex, and an echogenic agent; a and c are independently from 1 to
20; and b and d are independently vary from 1 to 100, and a
pharmaceutically acceptable carrier or excipient.

19. The composition of claim 18 wherein W^3 and X^3 are independently selected from the group consisting of $-C(CH_3)_2$, $-C((CH_2)_aOH)CH_3$, $-C((CH_2)_aOH)_2$, $-C((CH_2)_aCO_2H)CH_3$, $-C((CH_2)_aCO_2H)_2$, $-C((CH_2)_aNH_2)CH_3$, $C((CH_2)_aNH_2)_2$, $C((CH_2)_aNR^3R^4)_2$, $-NR^3$, and $-S$; Y^3 is
- 5 selected from the group consisting of $-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$, $-(CH_2)_a-NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$, $-(CH_2)_a-NR^3R^4$, and $-CH_2(CH_2OCH_2)_b-CH_2NR^3R^4$; Z^3 is selected from the group consisting of $-(CH_2)_a-CONH-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Dm$, $-(CH_2)_a-NHCO-Dm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Dm$, $-(CH_2)_a-$
- 10 NR^3R^4 , and $-CH_2(CH_2OCH_2)_b-CH_2NR^3R^4$; A_1 is a single or a double bond; B_1 , C_1 , and D_1 are independently selected from the group consisting of $-O-$, $-S-$, NR^3 , $(CH_2)_a-CR^1R^2$, and $-CR^1$; A_1 , B_1 , C_1 , and D_1 may together form a 6- to 10-membered carbocyclic ring or a 6- to 10-membered heterocyclic ring optionally containing one or more oxygen, nitrogen, or
- 15 sulfur atom; a_3 and b_3 are independently from 0 to 3; R^1 to R^4 , and R^{29} to R^{37} are independently selected from the group consisting of hydrogen, C_1 - C_{10} alkyl, C_5 - C_{12} aryl, C_1 - C_{10} alkoxy, C_1 - C_{10} polyhydroxyalkyl, C_5 - C_{12} polyhydroxyaryl, C_1 - C_{10} aminoalkyl, mono- or oligosaccharide, peptide with 2 to 30 amino acid units, $-CH_2(CH_2OCH_2)_b-CH_2-OH$, $-(CH_2)_a-CO_2H$,
- 20 $-(CH_2)_a-CONH-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-CONH-Bm$, $-(CH_2)_a-NHCO-Bm$, $-CH_2-(CH_2OCH_2)_b-CH_2-NHCO-Bm$, $-(CH_2)_a-OH$ and $-CH_2-(CH_2OCH_2)_b-CO_2H$; Bm and Dm are independently selected from the group consisting of a bioactive peptide containing 2 to 30 amino acid units, an antibody, a

mono- or oligosaccharide, a glycopeptide, a metal chelating agent, a
25 radioactive or nonradioactive metal complex, and an echogenic agent; a
and c are independently from 1 to 10; and b and d are independently from
1 to 30.

20. The composition of claim 19 wherein each of W^3 and X^3 is
 $C((CH_2)OH)_2$; Y^3 is $-(CH_2)_2-CONH-Bm$; Z^3 is $-(CH_2)_2-CONH-Dm$; A_1 is a
single bond; A_1 , B_1 , C_1 , and D_1 together form a 6-membered carbocyclic
ring; each of a_3 and b_3 is 1; R^{29} is galactose; each of R^{30} to R^{37} is
5 hydrogen; Bm is Octreotate; and Dm is bombesin (7-14).